

Aharonov-Bohm effect in curved space and cosmic strings

Yu.A.Sitenko A.V.Mishchenko

N.N.Bogolyubov Institute for Theoretical Physics

National Academy of Sciences of Ukraine

252143, Kiev, Ukraine

A quantum theory is developed for the scattering of a nonrelativistic particle in the field of a cosmic string regarded as a combination of a magnetic and gravitational strings. Allowance is made for the effects due to the finite transverse dimensions of the string under fairly general assumptions about the distribution of the magnetic field and spatial curvature in the string. It is shown that in a definite range of angles the differential cross section at all absolute values of the wave vector of the incident particle depends strongly on the magnetic flux of the string.

It is well known that in classical mechanics a complete description of electromagnetic effects can be made by means of the electromagnetic field strength, which acts directly on charged particles. In 1959, Aharonov and Bohm [1], using the Schrödinger equation, considered the scattering of an electron in an external static magnetic field produced by an infinitely long solenoid and found an effect that does not depend on the depth of penetration of the electrons into the region of magnetic force lines. This showed that in quantum mechanics the electromagnetic field acts on charged particles even in the case when the region in which the field is localized cannot be reached by the particles (see, for example, the reviews of Refs. [2] and [3]).

In this paper, we investigate the effect of curvature of space on the Aharonov-Bohm effect. Namely, we consider the situation in which there is not only a tube of magnetic force lines but also an external static cylindrically symmetric gravitational field with symmetry axis of that coincides with the axis of the magnetic tube. It is assumed, that at large distances from the symmetry axis the space becomes locally flat and the region of spatial curvature (the gravitational tube) may either coincide with the magnetic tube or include it or, finally, be included in it. We pose and solve the problem of the scattering of charged test particle in a space with gravitational and magnetic tubes.

It is appropriate to point out here that the simultaneous existence of magnetic and gravitational tubes is rather typical of models with spontaneous breaking of gauge symmetries. In such models, there arise vacuum structures in the form of strings (for example, Abrikosov-Nielsen-Olesen vortices [4],[5]), which are characterized, on the one hand, by a certain flux that one can reasonably call magnetic since it corresponds to the spontaneous breaking of a gauge degree of freedom and, on the other hand, by the value a condensate that spontaneously breaks a symmetry. To this condensate there corresponds a uniform distribution of mass along the string axis, and this is the source of the gravitational field of the type described above. We note that this latter field is rather weak and it is apparently for this reason that it has effectively escaped the attention of investigators. In particular, to the best of our knowledge the processes of scattering of particles by strings (vortices) usually take into account only the presence of the magnetic component, and the presence of the gravitational component is ignored (see, for example, Refs. [6] and [7]). It is to the correction of this, in our view, shortcoming that the present paper is devoted. We shall show that it is possible to take into account gravitational components of a much more general form than those inherent in strings in models with spontaneous breaking of gauge symmetries.

In what follows, we shall, for definiteness, consider strings, that are cosmological objects — so-called cosmic strings. According to modern ideas, the early stages in the evolution of the universe were characterized by high temperatures and a greater degree of symmetry than now, and as the universe cooled there was a series of phase

transitions with spontaneous symmetry breaking [8]. The topological defects formed as a result (monopoles, strings, and domain walls) may be stable, and the mere fact of their existence leads to many important consequences in cosmology [9]. Particular interesting cosmological objects are strings, which, in particular, can play the role of nucleation centers of galaxies and gravitational lenses [10].

We believe it is of interest to consider the situation in which there is flux of magnetic field along a gravitational string and to study the scattering of a test charged particle by such an object, in other words, to consider the standart (magnetic) Aharonov-Bohm effect in a Fierz space. This problem was actually posed for the first time in Ref. [11], in which some results were obtained; in particular attention was drawn to the difference between the scattering for spinor and scalar particles. In this paper, we make a more systematic and complete study of the problem. We determine the S - matrix for scattering in the case of combined singular magnetic and gravitational strings, establish the connection between the S - matrix and the scattering amplitude in this and in the more general case of nonsingular strings, and also take into account effects due to the finite transverse dimensions of the strings.

More precisely, we consider the spacetime with metric

$$ds^2 = -c^2 dt^2 + dz^2 + f(X, Y)(dX^2 + dY^2) \quad (1)$$

where $X = r \cos \phi$, $Y = r \sin \phi$ and f is the conformal factor of the metric of the surface $Z = \text{const.}$ This last surface has Gaussian curvature

$$K(X, Y) = -(2f)^{-1}(\partial_X^2 + \partial_Y^2) \ln f, \quad (2)$$

which satisfies the condition

$$K(X, Y) = 0 \quad (3)$$

for $r > r_K$. With regard to the closure of the region, $r \leq r_K$, we assume that here $K(X, Y)$ is a piecewise continuous function with singularities at isolated points or on isolated lines that are integrable with respect to the measure $f(X, Y)dXdY$. Then the total integrated curvature (in units of 2π) is given by

$$\Phi_K = \frac{1}{2\pi} \int_{r \leq r_K} dXdY f(X, Y) K(X, Y). \quad (4)$$

As we showed in Ref. [12], for $r > r_K$ (when condition (3) is satisfied) the conformal factor takes the form

$$F(X, Y) = \left(\frac{r}{r_0} \right)^{-2\Phi_K}. \quad (5)$$

Thus, taking into account (1), we obtain a relationship between the total curvature of the surface and the linear mass density of cosmic string

$$\Phi_K = 4GMc^{-2}. \quad (6)$$

In the case of a singular gravitational string, we have $r_K \rightarrow 0$ and

$$K(X, Y) = 2\pi\Phi_K \frac{\delta(X)\delta(Y)}{f(X, Y)} . \quad (7)$$

Along coordinate lines corresponding to the variable z , in the spacetime (1) a static magnetic field $B(X, Y)$ that satisfies the condition

$$B(X, Y) = 0 \quad (8)$$

for $r > r_B$ is directed. With regard to the closure of the region, $r \leq r_B$, we assume that here $B(X, Y)$ is a piecewise continuous function with singularities at isolated points or on isolated lines that are integrable with respect to the measure $f(X, Y)dXdY$. Then the total magnetic flux (in London units $2\pi\frac{\hbar c}{e}$) is given by

$$\Phi = \frac{e}{2\pi\hbar c} \int_{r \leq r_B} dXdY f(X, Y) B(X, Y) , \quad (9)$$

where e is the coupling constant of the matter to the gauge minus field ($-e$ is the charge of the test particle). In the case of a singular magnetic string, we have $r_B \rightarrow 0$ and

$$B(X, Y) = 2\pi\frac{\hbar c}{e}\Phi \frac{\delta(X)\delta(Y)}{f(X, Y)} . \quad (10)$$

In this paper, we study the quantum-mechanical scattering of nonrelativistic test particles by a cosmic string. Since the motion of the particles along the string axis z is free, we can make a restriction to considering the motion of particles on the surface $z = \text{const}$ (for more details see, for example, Ref. [11]). The Schrödinger equation for the wave function describing the stationary scattering state has the form

$$H\psi(X, Y) = \frac{\hbar^2 k^2}{2m} \psi(X, Y) , \quad (11)$$

where m and k are, respectively, the mass of the particle and the absolute value of its wave vector. Under the condition of axial symmetry of the magnetic field strength and the Gaussian curvature,

$$\partial_\phi B(X, Y) = 0, \quad \partial_\phi K(X, Y) = 0 , \quad (12)$$

we obtain for the Hamiltonian the expression

$$H = -\frac{\hbar^2}{2m} \left\{ \frac{1}{r^2 f} [(r\partial_r)^2 + (\partial_\phi - i\frac{e}{\hbar c}V + \frac{1}{2}i\sigma W)^2] + \sigma\frac{e}{\hbar c}B - \frac{1}{2}K \right\} , \quad (13)$$

where

$$V(r) = \int_0^r dr r f B, \quad W(r) = \int_0^r dr r f K , \quad (14)$$

$\sigma = \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ in the case of a spinor particle (ψ is a two-component column function) and $\sigma = 0$ in case of a scalar particle (ψ is a single-component function). As a radial variable, it is convenient to use the geodesic length in the radial direction:

$$\rho = \int_0^r dr f^{1/2}. \quad (15)$$

Then the metric (1) (under the condition of axial symmetry) takes the form

$$ds^2 = -c^2 dt^2 + dz^2 + d\rho^2 + \gamma^2 d\phi^2, \quad (16)$$

where

$$\gamma(\rho) = r(\rho) \sqrt{f[r(\rho)]}, \quad (17)$$

and the Hamiltonian (13) can be represented as follows:

$$H = -\frac{\hbar^2}{2m} \left\{ \gamma^{-1} \partial_\rho \gamma \partial_\rho + \gamma^{-2} [\partial_\phi - i \frac{e}{\hbar c} V + \frac{1}{2} i \sigma (1 - (\partial_\rho \gamma))]^2 + \right. \\ \left. + \sigma \frac{e}{\hbar c} \gamma^{-1} (\partial_\rho V) + \frac{1}{2} \gamma^{-1} (\partial_\rho^2 \gamma) \right\}. \quad (18)$$

The operator H acts in the space of functions with scalar product

$$(\psi, \psi') = \int_0^{2\pi} d\phi \int_0^\infty dr r f \psi^* \psi' = \int_0^{2\pi} d\phi \int_0^\infty d\rho \gamma \psi^* \psi'. \quad (19)$$

In contrast, the Hamiltonian in the absence of a string (i.e. with $\Phi = \Phi_K = 0$),

$$H_0 = -\frac{\hbar^2}{2m} (\partial_\rho^2 + \rho^{-1} \partial_\rho + \rho^{-2} \partial_\phi^2), \quad (20)$$

acts in the space of functions with scalar product

$$(\psi, \psi') = \int_0^{2\pi} d\phi \int_0^\infty d\rho \rho \psi^* \psi'. \quad (21)$$

We introduce a transformation of the operator and of the functions that leaves the form of equation (11) unchanged

$$\tilde{H} = \Xi H \Xi^{-1}, \quad \tilde{\psi} = \Xi \psi, \quad \tilde{H} \tilde{\psi} = \frac{\hbar^2 k^2}{2m} \tilde{\psi}. \quad (22)$$

Choosing

$$\Xi = \gamma^{1/2} \rho^{-1/2}, \quad (23)$$

we obtain for the transformed Hamiltonian the expression

$$\tilde{H} = -\frac{\hbar^2}{2m} \left\{ \partial_\rho^2 + \rho^{-1} \partial_\rho - \frac{1}{4} \rho^{-2} + \frac{1}{4} \gamma^{-2} (\partial_\rho \gamma)^2 + \right. \\ \left. + \gamma^{-2} \left[\partial_\phi - i \frac{e}{\hbar c} V + \frac{1}{2} i \sigma (1 - (\partial_\rho \gamma)) \right]^2 + \sigma \frac{e}{\hbar c} \gamma^{-1} (\partial_\rho V) \right\}; \quad (24)$$

then in the space of the transformed functions the scalar product is defined in accordance with (21), rather than (19).

We regard H_0 (20) as an unperturbed Hamiltonian and

$$\Delta H = \tilde{H} - H_0 \quad (25)$$

as an operator that describes a perturbing interaction, and we shall attempt to construct a scattering theory. Taking into account (20) and (24), we represent the operator (25) in the form

$$\Delta H = v(\vec{x}) + v^j(\vec{x})(-i\frac{\partial}{\partial x^j}) + v^{jj'}(\vec{x})(-\frac{\partial^2}{\partial x^j \partial x^{j'}}), \quad (26)$$

where we have introduced the notation $\vec{x} = (x^1, x^2)$, $x^1 = \rho \cos \phi$, $x^2 = \rho \sin \phi$ and $j, j' = 1, 2$.

If the coefficient functions v , v^j and $v^{jj'}$ decrease as $O(\rho^{-1-\epsilon})$ ($\epsilon > 0$) as $\rho \rightarrow \infty$, then in accordance with Ref. [13] the perturbation ΔH (26) has a short-range and scattering theory can be constructed in the usual manner (see, for example Refs. [14] and [15]). However, even for particle scattering by a purely magnetic string ($\Phi \neq 0$ and $\Phi_K = 0$) the perturbation ΔH has a long-range, since the coefficient function v^j decreases in the limit $\rho \rightarrow \infty$ as $O(\rho^{-1})$. Because of the long-range interaction, it is impossible to choose a plane wave as the incident wave, as noted by Aharonov and Bohm [1]. Nevertheless, in this case it is possible to develop a scattering theory and obtain in its framework the Aharonov-Bohm scattering amplitude (see Ref. [16]).

Hörmander [13] considered a class of perturbations of the form (26) containing both a short-range part and a long-range part characterized by real coefficient functions that decrease in the limit $\rho \rightarrow \infty$ as $O(\rho^{-\epsilon})$ ($0 < \epsilon \leq 1$), and he formulated certain additional requirements under which scattering theory can be developed. As he notes on page 417 of the translation of his monograph of Ref. [13] "the existence of modified wave operators is proved under the weakest sufficient conditions among all those known at the present time."

Hörmander's conditions are satisfied by the perturbation in the problem of scattering by a purely magnetic string,

$$v \sim O(\rho^{-2}) \text{ and } v^j \sim O(\rho^{-1}), \quad \rho \rightarrow \infty \quad (27)$$

(v and v^j are real functions, and $v^{jj'} = 0$), and, for example, perturbation in the problem of scattering by a Coulomb center,

$$v \sim O(\rho^{-1}), \quad \rho \rightarrow \infty \quad (28)$$

(v is a real function, and $v^j = v^{jj'} = 0$). In contrast, the perturbation in the problem of scattering by a cosmic string ($\Phi \neq 0$ and $\Phi_K \neq 0$) does not satisfy the Hörmander's conditions:

$$v \sim O(\rho^{-2}) \quad v^j \sim O(\rho^{-1}) \text{ and } v^{jj'} \sim O(1) \quad \rho \rightarrow \infty, \quad (29)$$

where v^j , in contrast to v and $v^{jj'}$ is a complex function (more precisely, imaginary part of v^j of order ρ^{-1} is due to the nondecrease of the real quantity $v^{jj'}$ in the limit $\rho \rightarrow \infty$). Nevertheless, even in this last case it is possible to develop a scattering theory, and in the present paper (see Ref. [17]) we shall construct wave operators explicitly.

In Sec. 2, we determine the S -matrix and the scattering in the case of a singular cosmic string ($r_B = 0$ and $r_K = 0$). In Sec. 3, we take into account the effects of the gravitational structure of the string (the finiteness of the transverse dimensions of the region of curvature of space, $r_K > 0$), and in Sec. 4 the effects of the magnetic structure of the string (finiteness of the transverse dimensions of the region of magnetic flux, $r_B > 0$). Sec. 5 is devoted to discussion of the results. Details of the derivation of the basic relations are given in Appendices A, B and C.

Acknowledgments

The work of Yu.A.S. and A.V.M. was supported by Swiss National Science Foundation Grant No. CEEC/NIS/96-98/7 IP 051219.

References

- [1] Y. Aharonov and D. Bohm, Phys. Rev. **115**, 485 (1959).
- [2] M. Peshkin and A. Tonomura, *The Aharonov-Bohm Effect (Lecture Notes in Physics*, Vol.340), Springer-Verlag, Berlin Heidelberg (1989).
- [3] G. N. Afanas'ev, Fiz. Elem. Chastits At. Yadra **21**, 172 (1990) [Sov. J. Part. Nucl. **21**, 74, (1990)].
- [4] A. A. Abrikosov, Zh. Eks. Teor. Fiz. **32**, 1442 (1957) [Sov. Phys. JETP **5**, 1174, (1957)].
- [5] H. B. Nielsen and P. Olesen, Nucl. Phys. B **61**, 45 (1973).
- [6] M. G. Alford, J. March-Russel, and F. Wilczek, Nucl. Phys. B **328**, 140 (1989).
- [7] C.-P. Ma, Phys. Rev. D **48**, 530 (1993).
- [8] D. A. Kirzhnits, Pis'ma Zh. Éksp. Teor. Fiz. **15**, 745 (1972) [JETP Lett. **15**, 539 (1972)]; D. A. Kirzhnits and A. D. Linde, Phys. Lett. B **42**, 471 (1972); D. A. Kirzhnits and A. D. Linde, Zh. Éksp. Teor. Fiz. **67**, 1263 (1974) [Sov. Phys. JETP **40**, 628 (1974)].
- [9] T. W. B. Kibble, J. Phys. A **9**, 1387 (1976); Phys. Rep. **67**, 183 (1980).
- [10] A. Vilenkin, Phys. Rev. D **23**, 852 (1981); Phys. Rev. D **24**, 2082 (1981).
- [11] Yu. A. Sitenko, Nucl. Phys. B **372**, 622 (1992).
- [12] A. V. Mishchenko and Yu. A. Sitenko, Yad. Fiz. **54**, 1158 (1991) [Sov. J. Nucl. Phys. **54**, 702 (1991)]; A. V. Mishchenko and Yu. A. Sitenko, Ann. Phys. (N.Y). **218**, 199 (1992).
- [13] L. Hörmander, *Analysis of Linear Partial Differential Operators IV*, Springer-Verlag, Berlin (1985)
- [14] A. G. Sitenko, *Scattering Theory* [Springer-Verlag, Berlin (1991)] 2nd ed., Vishcha Shkola, Kiev (1975).
- [15] M. C. Reed and B. Simon, *Methods of Modern Mathematical Physics*, Vol. 3 (Academic Press, New York, 1979)
- [16] S. N. M. Ruijsenaars, Ann. Phys. (N.Y). **146**, 1 (1983).
- [17] Yu. A. Sitenko and A. V. Mishchenko, JETP **81** (5), 831, (1995).